

Probiotec Limited 83 Cherry Lane, Laverton VIC 3026 Ph 1800 067 919 Fax +61 3 9369 6730 www.probiotec.com.au



Information Sheet - Feeding Guidelines for Palastart & Palastart Blue.



Recommended feeding rates for calves only. Prior to commencement of feeding to other animals please consult your local vet for feeding rates.

Palastart Milk Replacers are suitable for all types of animals.


Age of Animal	Volume - Twice Daily	Volume - Once Daily
*4 - 10 days	1.5 - 2.0 litres Add 125 - 150 grams of powder per litre.	2.0 - 2.5 litres Add 250 - 300 grams of powder per litre.
11 - 21 days	2.0 - 2.5 litres Add 125 - 150 grams of powder per litre.	2.5 - 3.0 litres Add 250 - 300 grams of powder per litre.
3 Weeks to Weaning	2.5 litres Add 125 - 150 grams of powder per litre.	3.0 litres Add 250 - 300 grams of powder per litre.

**Where possible colostrum and transition milk should be fed for the first four (4) days (See PBIS Colostrum feeding).*

Important.

- Palastart Blue milk replacer should be added at the higher addition rates (150 or 300 grams) shown in the table above.
- Palastart (Green) milk replacer should be added at the lower addition rates (125 or 250 grams) shown in the table above.
- Feeding guidelines are based on average weights for newborn calves.
- Smaller calves will require a smaller volume per feed.
- Large calves will require a larger volume per feed.
- Do not over feed.
- For twice a day feeding 8 - 9 hours should elapse between feeds.
- For once a day feeding feed at the same time each day. Probiotec recommend twice a day feeding for the first seven (7) days if possible.
- Feed the milk replacer at the same temperature each feeding.
- Changes to feeding volume and concentration should be made gradually.
- Clean fresh water should be available at all times.
- Good quality pellets or concentrate should be available ad lib from day Five (5).

While the suggestions and data contained herein are based on information believed to be reliable, NO WARRANTY, GUARANTEE, CONDITION OR REPRESENTATION IS GIVEN OR MADE IN RESPECT OF THE PRODUCTS OR IS TO BE TAKEN AS IMPLIED FROM ANY NEGOTIATIONS, CORRESPONDENCE, OR STATUTE AS TO THE QUALITY, MERCHANTABILITY OF FITNESS OF THE PRODUCTS FOR A PARTICULAR PURPOSE, except as set forth in Seller's Conditions of Sale. Purchasers should determine for themselves whether suggested formulations, example products or procedures are suitable for their own purposes. The specifications and suggested uses as may be described herein should not be construed to violate any patent or as warranty of non-infringement of any patent.



Probiotec Limited 83 Cherry Lane, Laverton VIC 3026 Ph 1800 067 919 Fax +61 3 9369 6730 www.probiotec.com.au



Information Sheet - Feeding Guidelines for Palastart & Palastart Blue.



Mixing.

- Mix the required quantity of powder (125 to 150 grams for twice a day and 250 to 300 grams for once a day) with one (1) litre of warm water to obtain the correct concentration for milk replacer.
- Fill the container with the required volume of clean warm water and slowly add the powder.
- Stir vigorously.
- Continue stirring until all of the powder and lumps have dissolved.
- Use as soon as possible after mixing.

For Further information on any of the points mentioned above contact your Palastart representative on 1800 067 919.

While the suggestions and data contained herein are based on information believed to be reliable, NO WARRANTY, GUARANTEE, CONDITION OR REPRESENTATION IS GIVEN OR MADE IN RESPECT OF THE PRODUCTS OR IS TO BE TAKEN AS IMPLIED FROM ANY NEGOTIATIONS, CORRESPONDENCE, OR STATUTE AS TO THE QUALITY, MERCHANTABILITY OF FITNESS OF THE PRODUCTS FOR A PARTICULAR PURPOSE, except as set forth in Seller's Conditions of Sale. Purchasers should determine for themselves whether suggested formulations, example products or procedures are suitable for their own purposes. The specifications and suggested uses as may be described herein should not be construed to violate any patent or as warranty of non-infringement of any patent.